

# WOKADINE INTRAUTERINE INFUSION<sup>®</sup>

An effective broad spectrum uterine treatment

## COMPOSITION:

Povidone Iodine B.P	10% w/v
(Available Iodine) B.P	1% w/v
Metronidazole B.P	1% w/v
Purified water B.P	q.s.

## INDICATIONS:

For treatment, prevention and control of infections related to;	
Infertility/repeat breeding	Pyometra
Metritis, Endometritis	Retention of placenta
Cervicitis and Vaginitis	Habitual/threatened abortion
Delayed uterine involution	Prolapse of uterus/vagina

## DOSAGE AND ADMINISTRATION

### Intrauterine use only

Withdraw required quantity of WOKADINE solution. Infuse into uterus with catheter and syringe. Treat once a day for 3 to 5 days as per need. Dose and number of doses must be based on condition / severity and therapeutic need assessed by the veterinarian.

Condition	Doses per day
Infertility and repeat breeding	20 – 30 ml dilute with equal amount of normal saline or sterile water for one day
Retention of placenta	30 – 60 ml for 2 -3 days
Pyometra, metritis	60 ml daily for 2 -3 days
Cervicitis and vaginitis	20 ml dilute with equal amount of sterile normal saline for 1 day

## PRESENTATION

60 ml sterile pouch with adaptor



Cow and calf



Healthy sheep and lamb



Retained placenta



## Osho Chemical Industries Limited

Industrial Area Office: Osho Complex, Sasio Rd. off Lunga lunga, Industrial Area  
 Postal Address: P. O. Box 49916 - GPO 00100 Nbi-Kenya Pilot Line: (+254) 020 391 2000, (+254) 0711-045 000,  
 Customer Service: (+254) 020 391 2500, (+254) 0711-045 500 Fax: (+254) 020 3912600  
 E-mail: oshochem@oshochem.com Website: www.oshochem.com