



## As tough | on wide range of pests



**Yenye nguvu dhidi ya aina mbalimbali ya wadudu waribifu shabani**

- **Broad spectrum.**
- **Systemic insecticide.**
- **With contact and residual action.**
- **Formulated and packaged for economic value.**

GUARANTEE | DHAMANA  
Acephate 75% w/w

Pilot lines: 0711 045000, 0732 167000, 020 3912000

4560 (SMS)

oshochem@oshochem.com

www.oshochem.com



**CHEMICAL INDUSTRIES LTD.**



ASATAF 75 SP is an organophosphorus broad spectrum systemic insecticide with contact and residual action. It has activity on chewing and sucking insects such as aphids, whiteflies, thrips and budworm. It can be applied as a foliar spray and for ground application.

ASATAF 75 SP ni dawa yenye nguvu za kuangamiza wadudu kwa njia ya mguso na kuenea kwenye mimea. Huweza pia kuendelea kuwauwa wadudu kwa muda mrefu. Huwauwa wadudu walao majani na wanaofyonza maji kwenye majani, kama vile aphids, thrips na budworm. Inaweza kunyunyizwa kwa matawi au mchanga.

DIRECTIONS FOR USE: Start spraying at first signs of insect attack, apply a volume of 1000 litres of water per hectare and ensure thorough and uniform plant coverage.

MAELEZO YA MATUMIZI: Anza kunyunyiza ASATAF 75 SP mara tu dalili za mashambulizi zinazopoonekana. Tumia kiwango cha lita 1000 za maji kwa hekta, na kuhakikisha mmea umelowa maji.

Precaution: The spray solution must be applied within two hours of its preparation. ASATAF 75 SP is non-phytotoxic when used at the recommended rates, however for usage in floriculture it is advisable to test varietal tolerance before large scale use.

Shauri: Mchanganyiko wa dawa lazima unyunyiziwe chini ya masaa mawili tangu unapotayarishwa. ASATAF 75 SP haina madhara kwenye mimea iwapo itatumika kwa kiwango kinachoshauriwa, lakini kwa unyunyizaji wa maua ni vyema kujaribu ustahimilivu wa aina mbali mbali za maua kabla ya matumizi makubwa.

CROP	PESTS	RATE	P.H.I
Vegetables e.g. French beans	Thrips, Aphids	10g/20L 500g/ha	3-7 days
Tomatoes	Caterpillars, Leaf miners, Whiteflies, Thrips, Aphids	10g/20L 500g/ha	3 days
Brassicas e.g. Cabbages etc	Cutworms, Spodoptera, Diamond-backmoth	10g/20L 500g/ha	7 days
Flowers e.g. Roses	Thrips, Aphids	50g in 100L water	3-7 days
Tobacco	Budworm, Leafminer, Aphids	40g in 20L water (soil applied)	15 days
Onions	Cutworm, Thrips	10g/20L 500g/ha	3-7 days
Cotton	Aphids	1kg/ha	15 days
Armyworms		10-15g/20L (0.5-0.75kg/ha)	



Aphid



Diamond-back moth



Spodoptera spp.



Thrip



Whitefly

Spray Interval: 14 days | Rudia baada ya siku 14.

SPRAY PREPARATION: Half fill the spray tank with the required volume of water, add the required amount of ASATAF 75 SP to the tank and stir. Add the remaining volume of water to fill the tank. Agitate or stir continuously until the powder is completely dissolved.

UCHANGANYISHAJI: Weka nusu ya kiwango kinachohitajika cha maji na kisha ongeza kipimo kinachostahili cha ASATAF 75 SP na kukoroga. Ongeza kiasi kilichobaki cha maji na kukoroga hadi dawa ichanganyike kikamilifu.

RE-ENTRY PERIOD: 24 hours after application for both man (without protective cover) and livestock.

KUINGIA TENA: Mahali palipo nyunyiziwa dawa hii hapafai kuingia binadamu (asiye na mavazi ya kinga) au wanyama, kabla ya masaa 24 kuisha.

Available packs | 30ml, 50ml, 100ml & 1Litre